

Massive Neutrinos, Lorentz Invariance Dominated Standard Model and the Phenomenological Approach to Neutrino Oscillations

(Phys. Scr.80 (2009) 025101)

Josip Šoln
Army Research Laboratory (ret.),
JZS Phys-Tech
Vienna, Virginia 22182, USA
soln.phystech@cox.net

August 13, 2009

Abstract

For the electroweak interactions, the massive neutrino perturbative kinematical procedure is developed in the massive neutrino Fock space; The perturbation expansion parameter is the ratio of neutrino mass to its energy. This procedure, within the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) modified electroweak Lagrangian, calculates the cross-sections with the new neutrino energy projection operators in the massive neutrino Fock space, resulting in the dominant Lorentz invariant Standard Model mass-less flavor neutrino cross-sections. As a consequence of the kinematical relations between the massive and massless neutrinos, some of the neutrino oscillation cross-sections are Lorentz invariance violating. But all these oscillating cross-sections, some of which violate the flavor conservation, being proportional to the squares of neutrino masses are practically unobservable in the laboratory. However, these neutrino oscillating cross-sections are consistent with the original Pontecorvo neutrino oscillating transition probability expression at short time (baseline), as presented by Dvornikov. From these comparisons, by mimicking the time dependence of the original Pontecorvo neutrino oscillating transition probability, one can formulate the dimensionless neutrino intensity-probability I , by phenomenologically extrapolating of the time t , or, equivalently the baseline distance L away from the collision point for the oscillating differential cross-section. For the incoming neutrino of 10MeV in energy and neutrino masses from Fritzsch analysis

with the neutrino mixing matrix of Harrison, Perkins and Scott, the baseline distances at the first two maxima of the neutrino intensity are $L \simeq 281km$ and $L \simeq 9279km$. The intensity I at the first maximum conserves the flavor, while at the second maximum, the intensities violate the flavor, respectively, in the final and initial state. At the end some details are given as to how these neutrino oscillations away from the collision point one should be able to verify experimentally. In material presented here one reinforces the notion that the mass-less flavor neutrino can be considered as the superposition of three massive neutrinos.

Introduction

The flavor changing neutrino oscillations experiments, such as, The Super-Kamiokande [1], SNO [2], KAMLAND [3] as well as Homestake Collaboration [4], clearly require massive neutrinos as have been exhibited, for example, by Bilenky, Giunti and Grimus [5], Giunti and Laveder [6] and Kayser [7]. In discussing the neutrino oscillations, one assumes that the left-handed flavor mass-less neutrino fields $\nu_{\alpha L}$, with $\alpha = e, \mu, \tau$, are unitary linear combinations of the massive neutrino fields ν_{iL} and analogously for the states (see [5-10]) and references therein),

$$\nu_{\alpha L} = U_{\alpha i} \nu_{iL}, |\nu_{\alpha}\rangle = U_{i\alpha}^{\dagger} |\nu_i\rangle \quad (i = 1, 2, 3; \alpha = e, \mu, \tau) \quad (1,2)$$

U is the unitary Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix and ν_{iL} is a left-handed neutrino field associated with mass m_i (see, for example [5-11]). In the study of flavor neutrino oscillations, the flavor state from (2) is used, via the Schrödinger equation, to calculate the oscillation probability $P_{\alpha\beta}(L)$ for the flavor neutrino oscillation transition $\nu_{\alpha} \longrightarrow \nu_{\beta}$ at a very large distance L ($\sim$ *thousands of km*) (see, for example [5-11]).

Not long after the suggestion by Pontecorvo [12] that massive neutrinos oscillate, which appears is the easiest to treat with the Schroedinger equation (consult [5-11]), people started looking as to how the neutrino masses influence the field theoretic treatment of the standard model (SM). One of the earliest paper that addressed that problem was by R. E. Schrock [13]. He studied mostly the decays where the neutrinos and antineutrinos appear in final states, such as the π , K and the nuclear β decays. Such decays are friendly to the application of the PMNS unitary transformations to the established SM. Here, the kinematics is then of the massive neutrinos augmented with the PMNS unitary matrix dependence. He proposed the tests that showed the promise for determination of neutrino masses and lepton mixing. Specifically, the proposed tests of the π and K decays could detect the neutrino masses in the 1 to 400 MeV range while the nuclear β decays in the 0.1 keV to 5 MeV range. Today it is known that all these ranges are too high as it appears that every neutrino mass is bellow 1 eV.

More recently, Li and Liu [14] have studied the connection of massive neutrinos to the SM by studying the inequivalent vaccua model (see [15] and references

therein); here, the transformation between the Fock space of neutrino mass states and the unitary inequivalent flavor states is a Bogoliubov transformation [14], [15]. This transformation, for instance, yields the corrections to the Pontecorvo neutrino oscillation probability that are of $O(m^2)$ (m denoting generically any of the three neutrino masses). However, the problem is that it also yields that the branching ratio of $W^+ \rightarrow e^+ + \nu_\mu$ to $W^+ \rightarrow e^+ + \nu_e$ is of $O(m^2)$ which contradicts the Hamiltonian that one started from. In other words, in the inequivalent vacuum model, there is a flavor changing current such as $W^+ \rightarrow e^+ + \nu_\mu$ and the branching ratio is different from that of the SM with zero neutrino masses. In Appendix, Li and Liu show that the neutrino oscillation effects are large enough to neglect the inequivalent vacuum model effects; that is, the sum of all three decay widths of $W^+ \rightarrow e^+ + \nu_{e,\mu,\tau}$ equals the width of $W^+ \rightarrow e^+ + \nu_e$ in the SM.

It is well known that all by itself, the standard model with mass-less flavor neutrinos has been remarkably successful in describing the laboratory experimental data, such as the neutrino scattering, at low and medium energies (see, for instance, M. Fukugita and T. Yanagida [8]; and C. Giunti and C. W. Kim [9]). So one can then ask whether the SM cross-sections can be derived when starting, instead with the mass-less flavor neutrino fields $\nu_{\alpha L}$, with the massive neutrino fields ν_{iL} . In this article the answer to this question is in affirmative.

To proceed in this direction, as suggested by (1), the application of the PMNS substitution rule (3) transforms the SM Lagrangian density with the mass-less neutrino fields into the one with the massive neutrino fields (4):

$$\text{The PMNS substitution : } \nu_{\alpha L} \rightarrow U_{\alpha i} \nu_{iL} \quad (3)$$

$$\begin{aligned} \alpha, \beta, \dots, \epsilon &= e, \mu, \tau; i, j, a, \dots, b = 1, 2, 3 : \\ l_{\alpha L} &= \begin{pmatrix} U_{\alpha i} \nu_{iL} \\ \alpha_L \end{pmatrix}, \epsilon_{L,R} = P_{L,R} \epsilon, P_{L,R} = \frac{1}{2} (1 \mp \gamma^5) \\ L_{W,int}^{Lepton} &= \frac{g}{\sqrt{2}} \sum_{\epsilon=e,\mu,\tau; i=1,2,3} [\bar{\nu}_{iL}(x) U_{i\epsilon}^\dagger \gamma^\mu \epsilon_L(x) W_\mu(x, +) \\ &\quad + \bar{\epsilon}_L(x) \gamma^\mu U_{\epsilon j} \nu_{jL}(x) W_\mu^\dagger(x, +)], \\ W^\mu(x, \pm) &= \frac{1}{\sqrt{2}} [W^\mu(x, 1) \mp i W^\mu(x, 2)] \\ L_{Z,int}^{Lepton} &= \frac{g}{c_W} Z_\mu(x) \sum_{\epsilon=e,\mu,\tau} \left[\bar{l}_{\epsilon L}(x) \frac{\tau_3}{2} \gamma^\mu l_{\epsilon L}(x) - s_W^2(-) \bar{\epsilon}(x) \gamma^\mu \epsilon(x) \right] \\ &= \frac{g}{4c_W} Z_\mu(x) \sum_{\epsilon=e,\mu,\tau, a,b=1,2,3} [\bar{\nu}_a(x) U_{a\epsilon}^\dagger \gamma^\mu (1 - \gamma^5) U_{\epsilon b} \nu_b(x) \\ &\quad + \bar{\epsilon}(x) \gamma^\mu [(4s_W^2 - 1) + \gamma^5] \epsilon(x)], \\ s_W &= \sin \theta_W, c_W = \cos \theta_W \end{aligned} \quad (4)$$

Since the Lagrangian densities(4) contains the massive neutrino fields, all the calculations are now done formally in the massive neutrino Fock space. The

mass-less neutrinos will be the mass state neutrinos in the limit of negligible masses as a result of the perturbative neutrino kinematical procedure.

Perturbative kinematical procedure for calculating the neutrino differential cross-sections

A free neutrino spinor field with the mass $m_i, i = 1, 2, 3$, is written generally with the creation and annihilation operators as

$$\begin{aligned}\nu_i(x) &= \frac{1}{(2\pi)^{\frac{3}{2}}} \int \frac{d^3q}{q^0} \sum_s e^{iqx} u(q, s) a(q, s) + e^{-iqx} v(q, s) b^\dagger(q, s) \\ q^0 &= (\vec{q}^2 + m_i^2)^{\frac{1}{2}}\end{aligned}\quad (5)$$

The perturbative kinematics is based on the fact that the neutrino mass $m_i (m_i < 1\text{eV})$ is generally much smaller than its absolute momentum value $|\vec{q}|$. Therefore it is convenient to start with the “mass-less” four-component neutrino momentum $q_{(\gamma)}^\mu$ with fixed flavor parameter γ

$$q_{(\gamma)}^\mu = (\vec{q}_{(\gamma)}, q_{(\gamma)}^0), q_{(\gamma)}^2 = 0, \gamma = e, \mu, \tau \quad (6)$$

Next, one assumes that under this flavor parameter γ are grouped together three massive neutrinos, say, ν_i with masses $m_i; i = 1, 2, 3$ then the difference among their energies $|\Delta q_{(i_1, i_2)}^0| = |(q_{(i_2)}^0 - q_{(i_1)}^0)| \cong |\Delta m_{i_2, i_1}^2 / q_{(\gamma)}^0| = |(m_{i_2}^2 - m_{i_1}^2) / q_{(\gamma)}^0|$ is much smaller than the quantum-mechanical uncertainty of the energy [16]. As a consequence, in this case with fixed γ it is impossible to distinguish the emission of neutrinos with different masses in the neutrino processes [16]. Hence, the three massive neutrinos, satisfying these quantum mechanical conditions, can be viewed as superposing themselves to form the flavor neutrino ν_γ [11, 16] as depicted by relations (1) and (2). With this in mind, with $q_{(i, \gamma)}^\mu = (\vec{q}_{(\gamma)}, (\vec{q}_{(\gamma)}^2 + m_i^2)^{\frac{1}{2}})$ as the four-momentum of the massive neutrino with mass m_i the perturbative kinematics can be presented as

$$\begin{aligned}q_{(i, \gamma)}^\mu &\simeq q_{(\gamma)}^\mu - g^{\mu 0} \frac{m_i^2}{2q_{(\gamma)}^0} + O(m_i^4), i = 1, 2, 3; \gamma(fixed) = (e, \mu, \tau); \\ \vec{q}_{(i, \gamma)} &= \vec{q}_{(\gamma)}, q_{(i, \gamma)}^0 \simeq q_{(\gamma)}^0 + \frac{m_i^2}{2q_{(\gamma)}^0} + O(m_i^4), \\ q_{(i, \gamma)}^2 &\simeq -m_i^2 + O(m_i^4)\end{aligned}\quad (7)$$

In (7) the terms with $O(m_i^4)$ will be neglected and the fixed parameter γ , as already established is the neutrino flavor. Thus with this perturbative kinematics $q_{(i, \gamma)}^\mu$ ceases to be a true Lorentz four-momentum, while $q_{(\gamma)}^\mu$ remains to be

so. However, since $q_{(\gamma)}^\mu$, as shown in (7), is the main part of $q_{(i,\gamma)}^\mu$, the main portions of cross-sections are expected to be Lorentz invariant (LI) while the Lorentz invariance violation (LIV) will be generated by $g^{\mu 0} m_i^2 / 2q_{(\gamma)}^0$ from (7). The LIV terms are expected to be very small due the smallness of neutrino masses. Taking these relations into account, within the massive neutrino Fock space the differential cross-sections with flavor neutrinos are calculated. The question, of course is: is the result consistent with the SM?

To continue, in analogy to $q_{(i,\gamma)}^\mu$, one now introduces $\hat{s}_{(i,\gamma)}$ and $s_{(i,\gamma)}$ to denote respectively, the helicity operators and eigenvalues for $i = 1, 2, 3$ massive neutrinos comprising the mass-less flavor neutrino ν_γ ; The helicity operator and eigenvalue of the mass-less flavor neutrino ν_γ are denoted, respectively, as $\hat{s}_{(\gamma)}$ and $s_{(\gamma)}$. And, the effects of the massive to mass-less-neutrino kinematical relation (7) on these helicity eigenvalues are simply, what one can call, the ordinary massive to mass-less neutrino helicity relation.

$$\begin{aligned}\hat{s}_{(i,\gamma)} &= \vec{q}_{(i,\gamma)} \cdot \vec{\sigma} / |\vec{q}_{(i,\gamma)}| = \vec{q}_{(\gamma)} \cdot \vec{\sigma} / |\vec{q}_{(\gamma)}| = \hat{s}_{(\gamma)}, \\ \hat{s}_{(i,\gamma)} &= \hat{s}_{(\gamma)} = \hat{s}_{(k,\gamma)} \implies s_{(i,\gamma)} = s_{(\gamma)} = s_{(k,\gamma)}, \text{ etc;} \\ i \text{ or } k, \dots &= 1, 2, 3; \gamma(\text{fixed}) = (e \text{ or } \mu \text{ or } \tau)\end{aligned}\tag{8}$$

As a consequence of (7) and (8), with spinor indices suppressed, the contractions of massive neutrino free-field operators with the massive neutrino and antineutrino states are, respectively

$$\begin{aligned}\langle 0 | \nu(x, l) | q_{(i,\gamma)}, s_{(i,\gamma)} \rangle &= \frac{1}{(2\pi)^{\frac{3}{2}}} e^{i(q_{(i,\gamma)} \cdot x)} \delta_{li} u(q_{(i,\gamma)}, s_{(i,\gamma)}), \\ \langle \bar{q}_{(j,\delta)}, s_{(j,\delta)} | \nu(x, k) | 0 \rangle &= \frac{1}{(2\pi)^{\frac{3}{2}}} e^{-i(q_{(j,\delta)} \cdot x)} \delta_{kj} v(q_{(j,\delta)}, s_{(j,\delta)})\end{aligned}\tag{9}$$

where $s_{(j,\delta)}$ and δ have the same kind interrelationship as $s_{(i,\gamma)}$ and γ in (8), etc.

Since, as shown in (7) to (9), the superposed three massive neutrinos contain the single flavor designation, either in the initial or final state, say, γ and δ , the process can be denoted as $\nu(\gamma) + \alpha(P_1) \rightarrow \nu(\delta) + \beta(P_2)$. From the Lagrangian densities (4) the amplitude and its Hermitian conjugate for the process containing massive neutrinos, are build around these respective flavor designations, γ and δ . so that the generic amplitudes are given, respectively, as

$$\begin{aligned}S_{amp} &\sim \sum_{i,j,\dots} \delta_4(q_{(i,\gamma)} + P_{(1)} - q_{(j,\delta)} - P_{(2)}) i M_{i,j,\dots}, \\ S_{amp}^\dagger &\sim \sum_{k,l,\dots} \delta_4(q_{(k,\gamma)} + P_{(1)} - q_{(l,\delta)} - P_{(2)}) (-i) M_{k,l,\dots}^*\end{aligned}\tag{10}$$

Here, the momenta indicate the actual massive neutrino-lepton scattering and different Latin indices indicate possibilities of summation with the U matrices

which, however, here is not necessary to be explicit. To derive the cross-section, with the help of (7), (8) and (9), one needs

$$\begin{aligned}
S_{amp}^* S_{amp} &= \sum_{i,j,\dots;k,l,\dots} \left\{ \delta_4^2(q_{(\gamma)} + P_{(1)} - q_{(\delta)} - P_{(2)}) \right. \\
&\quad + \delta_3^2(\vec{q}_{(\gamma)} + \vec{P}_{(1)} - \vec{q}_{(\delta)} - \vec{P}_{(2)}) \delta(q_{(\gamma)}^0 + P_{(1)}^0 - q_{(\delta)}^0 - P_{(2)}^0) \\
&\quad \times \delta'(q_{(\gamma)}^0 + P_{(1)}^0 - q_{(\delta)}^0 - P_{(2)}^0) \frac{1}{2} \left[\frac{m_i^2 + m_k^2}{q_{(\gamma)}^0} - \frac{m_j^2 + m_l^2}{q_{(\delta)}^0} \right] \\
&\quad \left. + O(m^4) \right\} (M_{i,j,\dots})(M_{k,l,\dots}^*) \\
&= \delta_4^2(q_{(\gamma)} + P_{(1)} - q_{(\delta)} - P_{(2)}) \sum_{i,j,\dots;k,l,\dots} (M_{i,j,\dots})(M_{k,l,\dots}^*) + O(m^4)
\end{aligned} \tag{11}$$

The final result in (11) is the consequence of general delta function property $\delta(x)\delta'(x) = 0$. The terms with $O(m^4)$, denoting the fourth power of products of variety of $m_i, m_k, \text{etc.}$, are neglected. It follows that while the Fock space contains the massive neutrino states, the cross-section will utilize the kinematics of massless flavor neutrinos.

Next, one needs the spinor expressions, appearing in (9), to reflect respectively, the kinematical and helicity relations in order to facilitate the cross-section calculations.

$$\begin{aligned}
u(q_{(i,\alpha)}, s_{(i,\alpha)}) &= \frac{m_i - \underline{q}_{(i,\alpha)}}{\sqrt{2(m_i + q_{(i,\alpha)}^0)}} u(m_i, \vec{0}, s_{(i,\alpha)}), \\
u(m_i, \vec{0}, s_{(i,\alpha)}) &= \pm 1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \\
\underline{q}_{(i,\alpha)} &= \gamma_\mu q_{(i,\alpha)}; \\
\bar{u}(q_{(i,\alpha)}, s_{(i,\alpha)}) &= \bar{u}(m_i, \vec{0}, s_{(i,\alpha)}) \frac{m_i - \underline{q}_{(i,\alpha)}}{\sqrt{2(m_i + q_{(i,\alpha)}^0)}}
\end{aligned} \tag{12}$$

For a process with γ and δ flavor designations, $\nu(\gamma) + \alpha(P_1) \longrightarrow \nu(\delta) + \beta(P_2)$, in cross-section evaluations, one will deal with the neutrino energy projection operator over the positive energy states. Furthermore, rather than averaging over, one simply sums over the massive neutrino helicity degrees of freedom. Consistent with the ordinary neutrino helicity relation (8), the sum is carried over only the equal helicity eigenvalues:

$$\begin{aligned}
s_{(i,\alpha)} &= s_{(k,\alpha)} = s_{(\alpha)} : \sum_{s_{(i,\alpha)}, s_{(k,\alpha)}} u(q_{(i,\alpha)}, s_{(i,\alpha)}) \otimes \bar{u}(q_{(k,\alpha)}, s_{(k,\alpha)}) \\
&= \sum_{s_{(\alpha)}} u(q_{(i,\alpha)}, s_{(\alpha)}) \otimes \bar{u}(q_{(k,\alpha)}, s_{(\alpha)}) \equiv \frac{1}{2} [q_{(i,\alpha)}, q_{(k,\alpha)}; +, c], \\
[q_{(i,\alpha)}, q_{(k,\alpha)}; +, c] &= \frac{(m_i - \underline{q}_{(i,\alpha)}) (1 + \gamma^0) (m_k - \underline{q}_{(k,\alpha)})}{2 \left[(m_i + q_{(i,\alpha)}^0) (m_k + q_{(k,\alpha)}^0) \right]^{\frac{1}{2}}}, \\
i \text{ and } k &= 1, 2, 3; \gamma = e \text{ or } \mu \text{ or } \tau
\end{aligned} \tag{13}$$

where the $+$ sign refers to the positive energy states and c refers to the fact that the equal helicity eigenvalues in the sum yield the coherent result. (The incoherent projection operators $[q_{(i,\alpha)}, q_{(k,\alpha)}; +, i]$ with unequal helicity eigenvalues $s_{(i,\alpha)} \neq s_{(k,\alpha)}$ are not dealt here.) The relation (13) defines the spinorial massive neutrino to mass-less neutrino helicity relation and it is consistent with the ordinary helicity relation (8). Carrying out the indicated operations in relation (13) as a power series over the neutrino masses /energy, one obtains for the neutrino energy projection operator over the positive energy states the following

$$\begin{aligned}
[q_{(i,\alpha)}, q_{(k,\alpha)}; +, c] &= \sum_{n=0}^2 [q_{(i,\alpha)}, q_{(k,\alpha)}; +, c]_n, \\
[q_{(i,\alpha)}, q_{(k,\alpha)}; +, c]_0 &= -\underline{q}_{(\alpha)}, \\
[q_{(i,\alpha)}, q_{(k,\alpha)}; +, c]_1 &= m_k + \frac{(m_k - m_i) \gamma^0 \underline{q}_{(\alpha)}}{2q_{(\alpha)}^0}, \\
[q_{(i,\alpha)}, q_{(k,\alpha)}; +, c]_2 &= -\frac{(m_k - m_i)^2 \underline{q}_{(\alpha)}}{8q_{(\alpha)}^{02}} + \frac{m_i m_k \gamma^0}{2q_{(\alpha)}^0}
\end{aligned} \tag{14}$$

The coherent energy operator $[q_{(i,\alpha)}, q_{(k,\alpha)}; +, c]$ generates the electroweak interactions that are the same as the SM interactions plus the LIV neutrino oscillation processes that are negligible since their cross-sections are proportional to the squares of neutrino masses and, as such, are essentially zero. Relation (14) is in essence the procedure for calculating the cross-sections for the processes requiring only the neutrino energy projection operators over the positive energy states.

Applications to the differential cross-section calculations

As established earlier and consistent with (11), the quasi-elastic electroweak process with massive neutrinos present, to $O(m^2)$, can be denoted with the

kinematics that uses just the mass-less flavor neutrinos.

$$\nu(q_{(\gamma)}) + \alpha(P_{(1)}) \longrightarrow \nu(q_{(\delta)}) + \beta(P_{(2)}); y = \frac{q_{(\gamma)}^0 - q_{(\delta)}^0}{q_{(\gamma)}^0} = \frac{P_{(2)}^0 - P_{(1)}^0}{q_{(\gamma)}^0} \quad (15)$$

where y is the normalized energy transfer. Now, although working in the massive neutrino Fock space, relation (11) says that the kinematics for the cross-sections for the quasi-elastic scattering of the massless flavor neutrinos is determined with flavor neutrino momenta according to $\delta_4(q_{(\gamma)} + P_{(1)} - q_{(\delta)} - P_{(2)})$. Furthermore, since (see also [8])

$$\int dy = \frac{1}{2\pi} \int d\sigma(q_{(\delta)}) d\sigma(P_{(2)}) \delta_4(q_{(\gamma)} + P_{(1)} - q_{(\delta)} - P_{(2)}), d\sigma(q) = \left(\frac{d^3q}{q^0} \right) \quad (16)$$

the normalized neutrino (or charged lepton) energy transfer $y = (q_{(\gamma)}^0 - q_{(\delta)}^0)/q_{(\gamma)}^0$ cannot affect Lorentz invariance of any of the differential cross-sections.

Also, in view of (11), the cross-section normalization factor is defined with respect to the massless flavor neutrino momenta.

$$B = \frac{1}{(2\pi)^6} \left| (P_{(1)} \cdot q_{(\gamma)})^2 - P_{(1)}^2 q_{(\gamma)}^2 \right|^{\frac{1}{2}} = \frac{1}{(2\pi)^6} |(P_{(1)} \cdot q_{(\gamma)})|$$

In explicit evaluations, one uses the following short-hand notations:

$$m_{\alpha\beta} = \sum_i U_{\alpha i} m_i U_{i\beta}^\dagger, m_{\alpha\beta}^2 = \sum_i U_{\alpha i} m_i^2 U_{i\beta}^\dagger \quad (17)$$

Deriving the differential cross-sections with new energy projection operators for the flavor neutrino processes within the massive neutrino Fock space—

$\frac{d\sigma_W}{dy}$ —From the Lagrangian density in (4), the free neutrino field (5), the kinematical relation (7), the relations (8) and (9), one derives in the usual way the W -exchange S_W and $S_W^\dagger$ matrix elements for the process in (15). Specifically, with the Fierz rearrangement and repeated indices summing up, one has,

$$\begin{aligned} S_W &= \sum_{i,j} \delta_4(q_{(i,\gamma)} + P_{(1)} - q_{(j,\delta)} - P_{(2)}) \delta_{\alpha\beta} U_{\delta j} U_{i\gamma}^\dagger U_{j\beta}^\dagger U_{\alpha i} \frac{ig^2}{(2\pi)^2 8M_W^2} \\ &\quad \times \bar{u}(q_{(j,\delta)}, s_{(j,\delta)}) \gamma^\mu (1 - \gamma^5) u(q_{(i,\gamma)}, s_{(i,\gamma)}) \bar{u}(P_{(2)}, r_2) \gamma_\mu u(P_{(1)}, r_1) \end{aligned} \quad (18)$$

and $S_W^\dagger$ is obtained from (18) as shown in (10). The contribution to the process (15) due to the W -exchange from (18), after taking into account (11),

(17), $\sqrt{2}g^2 = 8M_W^2 G$, and the fact that $s_{(i,\gamma)}, s_{(j,\delta)}, \dots$, obey, respectively, the ordinary and spinorial helicity relation, (8) and (13), the standard procedure gives,

$$\begin{aligned}
s_{(i,\gamma)} &= s_{(g,\gamma)} = s_{(\gamma)}; s_{(h,\delta)} = s_{(j,\delta)} = s_{(\delta)} : \\
\frac{d\sigma_W(m)}{dy} &= \frac{d\sigma_W^{(c,c)}(m)}{dy} = \frac{G^2}{4\pi |(P_{(1)} \cdot q_{(\gamma)})|} \frac{\delta_{\alpha\beta}}{2^5} \sum_{i,j;g,h} \left(U_{h\delta}^\dagger U_{\gamma g} U_{\beta h} U_{g\alpha}^\dagger \right) \left(U_{\delta j} U_{i\gamma}^\dagger U_{j\beta}^\dagger U_{\alpha i} \right) \\
&\times \left[Tr \left(M_1 - \underline{P}_{(1)} \right) \gamma_\nu (1 - \gamma^5) \left(M_2 - \underline{P}_{(2)} \right) \gamma_\mu (1 - \gamma^5) \right] \\
&\times \left[Tr \left[q_{(i,\gamma)}, q_{(g,\gamma)}; +, c \right] \gamma^\nu (1 - \gamma^5) \left[q_{(h,\delta)}, q_{(j,\delta)}; +, c \right] \gamma^\mu (1 - \gamma^5) \right]
\end{aligned} \tag{19}$$

where m symbolically denotes dependence on $m_{1,2,3}$. Next, the coherent energy operator expansion according to (14), with gamma matrices traces carried out, yields

$$\begin{aligned}
\frac{d\sigma_W(m)}{dy} &= \frac{d\sigma_W(SM)}{dy} \left[1 + \frac{m_{\alpha\alpha}^2}{4} \left(\frac{1}{q_{(\gamma)}^2} + \frac{1}{q_{(\delta)}^2} \right) \right] - \frac{2G^2 \delta_{\alpha\beta}}{\pi |(P_{(1)} \cdot q_{(\gamma)})|} \\
&\times \left\{ \frac{\delta_{\alpha\gamma}}{4} m_{\alpha\delta} m_{\delta\alpha} \left[\frac{(P_{(1)} \cdot q_{(\gamma)}) (P_{(2)} \cdot q_{(\delta)})}{q_{(\delta)}^2} + \frac{2P_{(2)}^0 (P_{(1)} \cdot q_{(\gamma)})}{q_{(\delta)}^0} \right] \right. \\
&\left. + \frac{\delta_{\beta\delta}}{4} m_{\alpha\gamma} m_{\gamma\alpha} \left[\frac{(P_{(1)} \cdot q_{(\gamma)}) (P_{(2)} \cdot q_{(\delta)})}{q_{(\gamma)}^2} + \frac{2P_{(1)}^0 (P_{(2)} \cdot q_{(\delta)})}{q_{(\gamma)}^0} \right] \right\} + O(m^4), \\
\frac{d\sigma_W(SM)}{dy} &= \frac{2G^2 \delta_{\alpha\beta} \delta_{\alpha\gamma} \delta_{\beta\delta}}{\pi |(P_{(1)} \cdot q_{(\gamma)})|} (P_{(1)} \cdot q_{(\gamma)}) (P_{(2)} \cdot q_{(\delta)})
\end{aligned} \tag{20}$$

One can notice that, while the negligible LIV is associated with the neutrino mass, the LI Standard Model result is formally identified with zero neutrino mass limits

$$\frac{d\sigma_W(m)}{dy} = \frac{d\sigma_W(SM)}{dy} + O(m^2; LIV) \tag{21}$$

one can summarize the neutrino flavor transitions for the W -exchange neutrino processes. Flavor conserving are: LI to $O(m=0)$ terms and negligible LIV to $O(m^2)$ terms. Flavor violating are: negligible LIV to $O(m^2)$ terms.

$\frac{d\sigma_Z}{dy}$ —As in the previous case, from the Lagrangian density in (4), the free neutrino field (5), the kinematical relation (7), the contractions (9), one derives in the usual way the Z -exchange S_Z and $S_Z^\dagger$ matrix elements for the process in (15). Specifically, one has

$$\begin{aligned}
S_Z &= \sum_{i,j} \delta_4 (q_{(i,\gamma)} + P_{(1)} - q_{(j,\delta)} - P_{(2)}) \delta_{ij} \delta_{\alpha\beta} U_{\delta j} U_{i\gamma}^\dagger \\
&\times \frac{ig^2}{(2\pi)^2 16c_W^2 M_Z^2} [\bar{u}(q_{(j,\delta)}, s_{(j,\delta)}) \gamma^\mu (1 - \gamma^5) u(q_{(i,\gamma)}, s_{(i,\gamma)})] \\
&\times \bar{u}(P_{(2)}, r_2) \gamma_\mu (4s_W^2 - 1 + \gamma^5) u(P_{(1)}, r_1)
\end{aligned} \quad (22)$$

while $S_Z^\dagger$ is obtained from the S_Z through the Hermitian conjugation. In what follows, one will find the following shorthand notation very useful:

$$\begin{aligned}
w_0 &= s_W^2, w_1 = 2s_W^2 - 1, z_1 = s_W^2 (2s_W^2 - 1) + \frac{1}{4}, z_2 = s_W^2 (2s_W^2 - 1), \\
z_3 &= s_W^2 - \frac{1}{4}, z_4 = s_W^2 (s_W^2 - 1) + \frac{1}{4}, z_1 + z_3 = 2s_W^4
\end{aligned} \quad (23)$$

After taking into account that $c_W^2 M_Z^2 = M_W^2$ and the fact that the helicities, $s_{(i,\gamma)}, s_{(j,\delta)}, \dots$, obey both the ordinary and the spinorial helicity relations (8) and (13), the standard procedure yields the general expressions:

$$\begin{aligned}
s_{(i,\gamma)} &= s_{(k,\gamma)} = s_{(\gamma)}; s_{(l,\delta)} = s_{(j,\delta)} = s_{(\delta)} : \\
\frac{d\sigma_Z(m)}{dy} &= \frac{d\sigma_Z^{(c,c)}(m)}{dy} = \frac{G^2}{4\pi |(P_{(1)}, q_{(\gamma)})|} \frac{\delta_{\alpha\beta}}{2^5} \sum_{i,j;k,l} \left(U_{\delta j} U_{i\gamma}^\dagger U_{l\delta}^\dagger U_{\gamma k} \right) \delta_{ij} \delta_{kl} \\
&\times \left[Tr \left(M_1 - \underline{P}_{(1)} \right) \gamma_\nu \left(2z_3 + \frac{1}{2} \gamma^5 \right) \left(M_2 - \underline{P}_{(2)} \right) \gamma_\mu \left(2z_3 + \frac{1}{2} \gamma^5 \right) \right] \\
&\times \left[Tr [q_{(i,\gamma)}, q_{(k,\gamma)}; +, c] \gamma^\nu (1 - \gamma^5) [q_{(l,\delta)}, q_{(j,\delta)}; +, c] \gamma^\mu (1 - \gamma^5) \right]
\end{aligned} \quad (24)$$

The coherent energy operator expansion according to (14), with evaluating the traces of gamma matrices, yields

$$\begin{aligned}
\frac{d\sigma_Z(m)}{dy} &= \frac{d\sigma_Z(SM)}{dy} \left[1 + \frac{m_{\gamma\gamma}^2}{4} \left(\frac{1}{q_{(\gamma)}^2} + \frac{1}{q_{(\delta)}^2} \right) \right] - \frac{G^2 m_{\gamma\delta} m_{\delta\gamma} \delta_{\alpha\beta}}{8\pi |(P_{(1)} \cdot q_{(\gamma)})|} \\
&\times \left\{ 2 \left(\frac{1}{q_{(\gamma)}^2} + \frac{1}{q_{(\delta)}^2} \right) [M_1 M_2 z_2 (q_{(\gamma)} \cdot q_{(\delta)}) \right. \\
&+ (P_{(1)} \cdot q_{(\delta)}) (P_{(2)} \cdot q_{(\gamma)}) (z_1 + z_3) \\
&+ (P_{(1)} \cdot q_{(\gamma)}) (P_{(2)} \cdot q_{(\delta)}) (z_1 - z_3)] \\
&+ \frac{4}{q_{(\delta)}^0} [M_1 M_2 z_2 q_{(\gamma)}^0 + P_{(1)}^0 (P_{(2)} \cdot q_{(\gamma)}) (z_1 + z_3) \\
&+ P_{(2)}^0 (P_{(1)} \cdot q_{(\gamma)}) (z_1 - z_3)] + \frac{4}{q_{(\gamma)}^0} [M_1 M_2 z_2 q_{(\delta)}^0 \\
&+ P_{(2)}^0 (P_{(1)} \cdot q_{(\delta)}) (z_1 + z_3) + P_{(1)}^0 (P_{(2)} \cdot q_{(\delta)}) (z_1 - z_3)] \Big\} \\
&+ O(m^4) \\
\frac{d\sigma_Z(SM)}{dy} &= \frac{G^2 \delta_{\alpha\beta} \delta_{\gamma\delta}}{\pi |(P_{(1)} \cdot q_{(\gamma)})|} \\
&\times [M_1 M_2 z_2 (q_{(\gamma)} \cdot q_{(\delta)}) + (P_{(1)} \cdot q_{(\delta)}) (P_{(2)} \cdot q_{(\gamma)}) (z_1 + z_3) \\
&+ (P_{(1)} \cdot q_{(\gamma)}) (P_{(2)} \cdot q_{(\delta)}) (z_1 - z_3)]
\end{aligned} \tag{25}$$

Here also, the negligible LIV is associated with the neutrino mass while the LI Standard Model result is identified with formally zero neutrino mass limits :

$$\frac{d\sigma_Z(m)}{dy} = \frac{d\sigma_Z(SM)}{dy} + O(m^2; LIV), \tag{26}$$

While the terms of $O(m=0)$ are LI and flavor conserving, the negligible LIV terms of $O(m^2)$ are either flavor violating or flavor conserving.

$\frac{d\sigma_{\{W,Z\}}}{dy}$ —Here, the differential cross-section for the quasi-elastic neutrino scattering (15) due to the overlapping S —*matrix* elements from the W — and Z —is given as a sum of its components after taking into account relations (11), (18), and (22). Importantly, again taking into account the fact that helicities, $s_{(i,\gamma)}, s_{(j,\delta)}, \dots$, obey both the ordinary and spinorial helicity relations, (8) and (13), the standard procedure yields the general expression

$$\begin{aligned}
s_{(e,\gamma)} &= s_{(g,\gamma)} = s_{(i,\gamma)} = s_{(k,\gamma)} = s_{(\gamma)}; \\
s_{(h,\delta)} &= s_{(f,\delta)} = s_{(l,\delta)} = s_{(j,\delta)} = s_{(\delta)}; \\
\frac{d\sigma_{\{W,Z\}}(m)}{dy} &= \frac{d\sigma_{\{W,Z\}}^{(c,c)}(m)}{dy} = \frac{G^2}{8\pi |(P_{(1)}, q_{(\gamma)})|} \frac{\delta_{\alpha\beta}}{2^5} \\
&\times \left\{ \left[Tr \left(M_1 - \underline{P}_{(1)} \right) \gamma_\nu (4z_3 + \gamma^5) \left(M_2 - \underline{P}_{(2)} \right) \gamma_\mu (1 - \gamma^5) \right] \right. \\
&\times \left[\sum_{g,h;e,f} \left(U_{h\delta}^\dagger U_{\gamma g} U_{\beta h} U_{g\alpha}^\dagger U_{\delta f} U_{e\gamma}^\dagger \delta_{ef} \right) \right. \\
&\times Tr \left[q_{(e,\gamma)}, q_{(g,\gamma)}; +, c \right] \gamma^\nu (1 - \gamma^5) \left[q_{(h,\delta)}, q_{(f,\delta)}; +, c \right] \gamma^\mu (1 - \gamma^5) \left. \right] \\
&+ \left[\sum_{k,l;i,j} \left(U_{l\delta}^\dagger U_{\gamma k} U_{\delta j} U_{i\gamma}^\dagger U_{\alpha i} U_{j\beta}^\dagger \delta_{kl} \right) \right. \\
&\times Tr \left[q_{(i,\gamma)}, q_{(k,\gamma)}; +, c \right] \gamma^\nu (1 - \gamma^5) \left[q_{(l,\delta)}, q_{(j,\delta)}; +, c \right] \gamma^\mu (1 - \gamma^5) \left. \right] \left. \right\} \quad (27)
\end{aligned}$$

where one took into account the identity:

$$\begin{aligned}
&Tr \left(M_1 - \underline{P}_{(1)} \right) \gamma_\nu (4z_3 + \gamma^5) \left(M_2 - \underline{P}_{(2)} \right) \gamma_\mu (1 - \gamma^5) \\
&= Tr \left(M_1 - \underline{P}_{(1)} \right) \gamma_\nu (1 - \gamma^5) \left(M_2 - \underline{P}_{(2)} \right) \gamma_\mu (4z_3 + \gamma^5)
\end{aligned}$$

Of course, one cannot avoid the coherent energy operator expansion according to (14), and evaluating the traces of gamma matrices one obtains

$$\begin{aligned}
\frac{d\sigma_{\{W,Z\}}(m)}{dy} &= \frac{d\sigma_{\{W,Z\}}(SM)}{dy} \left[1 + \frac{m_{\gamma\gamma}^2}{4} \left(\frac{1}{q_{(\gamma)}^2} + \frac{1}{q_{(\delta)}^2} \right) \right] - \frac{G^2 m_{\gamma\delta} m_{\delta\gamma} \delta_{\alpha\beta}}{2\pi |(P_{(1)}, q_{(\gamma)})|} \\
&\times \left\{ \left[M_1 M_2 w_0 (q_{(\gamma)} \cdot q_{(\delta)}) + w_1 (P_{(1)}, q_{(\gamma)}) (P_{(2)}, q_{(\delta)}) \right] \right. \\
&\times \left(\frac{\delta_{\alpha\gamma}}{q_{(\gamma)}^2} + \frac{\delta_{\beta\delta}}{q_{(\delta)}^2} \right) + \frac{2\delta_{\alpha\gamma}}{q_{(\delta)}^0} \left[M_1 M_2 w_0 q_{(\gamma)}^0 + w_1 (P_{(1)}, q_{(\gamma)}) P_{(2)}^0 \right] \\
&+ \frac{2\delta_{\beta\delta}}{q_{(\gamma)}^0} \left[M_1 M_2 w_0 q_{(\delta)}^0 + w_1 (P_{(2)}, q_{(\delta)}) P_{(1)}^0 \right] \left. \right\} + O(m^4), \\
\frac{d\sigma_{\{W,Z\}}(SM)}{dy} &= \frac{2G^2 \delta_{\alpha\beta} \delta_{\alpha\gamma} \delta_{\gamma\delta}}{\pi |(P_{(1)}, q_{(\gamma)})|} \left[M_1 M_2 w_0 (q_{(\gamma)} \cdot q_{(\delta)}) + w_1 (P_{(1)}, q_{(\gamma)}) (P_{(2)}, q_{(\delta)}) \right] \quad (28)
\end{aligned}$$

Again, the negligible LIV is associated with the neutrino mass while the LI Standard Model result is identified with formally zero neutrino mass limits :

$$\frac{d\sigma_{\{W,Z\}}(m)}{dy} = \frac{d\sigma_{\{W,Z\}}(SM)}{dy} + O(m^2; LIV) \quad (29)$$

The overlapping W - and Z - exchanges cross-section terms of $O(m=0)$ are LI and flavor conserving, while the negligible terms of $O(m^2)$ carry the LIV terms with, both, the conserved and violated flavor.

Phenomenological neutrino cross-section intensity from time extrapolation of the neutrino oscillation scattering

Dvornikov in the classical field theoretical model [17] quotes the neutrino oscillation-transition probability with the characteristic Pontecorvo dimensionless argument $\Delta m^2 t / 4E$,

$$P(t) = \sin^2(2\Theta_{Vac}) \sin\left(\frac{\Delta m^2 t}{4E}\right) \quad (30)$$

where Θ_{Vac} is the vacuum mixing angle, $\Delta m^2 = m_1^2 - m_2^2$ is the mass squared difference and E is the energy of the system (detailed description of these and other parameters is in [17]). The interest in [17] comes from the fact that at $t = E^{-1}$ one can expand (3) for small Pontecorvo argument as $\sin\left(\frac{\Delta m^2}{4E^2}\right) = \frac{\Delta m^2}{4E^2} + \dots$. Or, turning the logic around, one could think of the scattering theory presented here as an approximation of a more general description, yet to be derived, where $\sin\left(\frac{\Delta m^2}{4E^2}\right)$ occurred but for the laboratory measurements was approximated with $\frac{\Delta m^2}{4E^2}$. So, with this reversed logic, one can assume that the calculated neutrino intensity, defined appropriately from differential cross-sections, can be analyzed at the practical baseline region if in it every dimensionless Pontecorvo argument of the type $\frac{\Delta m^2}{4E^2}$ is replaced with $\sin\left(\frac{\Delta m^2 t}{4E}\right)$. Other terms remain unaffected and, if proportional to the squares of neutrino masses, may even be neglected. However, the recoil, which Dvornikov [17] in the classical approach did not have, here will have to be taken into account and, if possible, averaged out in the calculated neutrino intensity in the short baseline region.

To pursue the idea from reference [17] on the phenomenological level, say, on the neutrino scattering process as described by the differential cross-section (28), it is necessary to work in the laboratory frame, $P_{(1)} = (\vec{0}, M_1)$. Next thing is to concentrate on the incoming neutrino energy in (28) with the help of the neutrino energy transfer

$$\begin{aligned} y &= \frac{q_{(\gamma)}^0 - q_{(\delta)}^0}{q_{(\gamma)}^0} = \frac{P_{(2)}^0 - M_1}{q_{(\gamma)}^0} \\ q_{(\delta)}^0 &= (1 - y) q_{(\gamma)}^0 \end{aligned} \quad (31)$$

So that after substituting (31) into (28) one obtains for the differential cross-section in (28)

$$\begin{aligned}
\frac{d\sigma_{\{W,Z\}}(m)}{dy} &= \frac{d\bar{\sigma}_{\{W,Z\}}(SM)}{dy} \delta_{\alpha\beta} \left\{ \delta_{\alpha\gamma} \delta_{\gamma\delta} + \delta_{\alpha\gamma} \delta_{\gamma\delta} \frac{m_{\gamma\gamma}^2}{4q_{(\gamma)}^2} \left[1 + \frac{1}{(1-y)^2} \right] \right. \\
&\quad \left. - \frac{m_{\gamma\delta} m_{\delta\gamma}}{4q_{(\gamma)}^2} \left[\delta_{\alpha\gamma} + \frac{\delta_{\beta\delta}}{(1-y)^2} \right] \right\} \\
&\quad - \frac{G^2 m_{\gamma\delta} m_{\delta\gamma} \delta_{\alpha\beta}}{\pi |(P_{(1)} \cdot q_{(\gamma)})|} \left\{ w_0 M_1 M_2 \left(\frac{\delta_{\alpha\gamma}}{(1-y)} + \delta_{\beta\delta} (1-y) \right) \right. \\
&\quad \left. + w_1 \left(-\frac{\delta_{\alpha\gamma} M_1 P_{(2)}^0}{(1-y)} + \frac{\delta_{\beta\delta} (P_{(2)} \cdot q_{(\delta)}) M_1}{q_{(\gamma)}^0} \right) \right\}
\end{aligned} \tag{32}$$

$$\begin{aligned}
\frac{d\bar{\sigma}_{\{W,Z\}}(SM)}{dy} &= \frac{2G^2}{\pi |(P_{(1)} \cdot q_{(\gamma)})|} [M_1 M_2 w_0 (q_{(\gamma)} \cdot q_{(\delta)}) \\
&\quad - w_1 M_1 q_{(\gamma)}^0 (P_{(2)} \cdot q_{(\delta)})]
\end{aligned} \tag{33}$$

In order to bring out the new features, one "removes" the SM cross-section from the newly derived cross-section, through the definition of the neutrino cross-section intensity which, as suggested by (30), will allow the introduction of the dimensionless Pontecorvo argument (30).

$$\begin{aligned}
I_{\{W,Z\}}(m, \theta) &= \left[\frac{d\sigma_{\{W,Z\}}(m)}{dy} - \frac{d\bar{\sigma}_{\{W,Z\}}(SM)}{dy} \right] \left[\frac{d\bar{\sigma}_{\{W,Z\}}(SM)}{dy} \right]^{-1} \\
&= \delta_{\alpha\beta} \left\{ \delta_{\alpha\gamma} \delta_{\gamma\delta} \frac{m_{\gamma\gamma}^2}{4q_{(\gamma)}^2} \left[1 + \frac{1}{(1-y)^2} \right] \right. \\
&\quad \left. - \frac{m_{\gamma\delta} m_{\delta\gamma}}{4q_{(\gamma)}^2} \left[\delta_{\alpha\gamma} + \frac{\delta_{\beta\delta}}{(1-y)^2} \right] \right\} \\
&\quad - \frac{m_{\gamma\delta} m_{\delta\gamma} \delta_{\alpha\beta}}{2} \left\{ \left[w_0 M_1 M_2 \left(\frac{\delta_{\alpha\gamma}}{(1-y)} + \delta_{\beta\delta} (1-y) \right) \right. \right. \\
&\quad \left. \left. + w_1 \left(-\frac{\delta_{\alpha\gamma} M_1 P_{(2)}^0}{(1-y)} + \frac{\delta_{\beta\delta} (P_{(2)} \cdot q_{(\delta)}) M_1}{q_{(\gamma)}^0} \right) \right] \right. \\
&\quad \left. \times \left[w_0 M_1 M_2 (q_{(\gamma)} \cdot q_{(\delta)}) - w_1 M_1 q_{(\gamma)}^0 (P_{(2)} \cdot q_{(\delta)}) \right]^{-1} \right\}
\end{aligned} \tag{34}$$

$$\vec{q}_{(\gamma)} \cdot \vec{P}_{(2)} = q_{(\gamma)}^0 |\vec{P}_{(2)}| \cos \theta, \quad y = \frac{2M_1 q_{(\gamma)}^0 \cos^2 \theta}{(M_1 + q_{(\gamma)}^0)^2 - q_{(\gamma)}^2 \cos^2 \theta} \tag{35}$$

Where the dimensionless intensity $I_{\{W,Z\}}(m, \theta)$ from (34) will serve also as unnormalized probability. As shown in (31) and (35), the angle θ -dependence

comes through the kinematics. The expression for the normalized neutrino energy transfer y follows from scattering kinematics and the energy-momentum conservation (more details can be found also in [9]). As seen in (35), for fixed $q_{(\gamma)}^0$, y depends only on $\cos^2 \theta$ which could be averaged over straightforwardly.

The relation (34), of course, is an instantaneous intensity at the time of interaction. This instant can be defined as $t_0 = 1/q_{(\gamma)}^0$. For $q_{(\gamma)}^0 = 10 \text{ MeV}$, $t_0 = 6.58 \times 10^{-23} \text{ sec.}$; one can also define a "distance" with $L_0 = 1/q_{(\gamma)}^0$ to give for the same neutrino energy $L_0 = 1.97 \times 10^{-12} \text{ cm}$. Of course, what counts are relative times and distances. Here, t_0 and L_0 are given for the sake of convenience. Now, as is described at the beginning of this section, one can perform the time extrapolation with sinus functions on the arguments that are in the Pontecorvo forms while others, with neutrino mass square dependences, can be dropped. The result of this procedure is, where for the sake of simplicity, the processes are denoted with just the flavor quantum numbers. One has

$$\begin{aligned}
I_{\{W,Z\}}(m, \theta) &= I_1(P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \\
&\quad + I_2(P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \\
&\quad + I_3(P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}), \\
I_1(P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) &= \delta_{\alpha\beta} \delta_{\alpha\gamma} \delta_{\gamma\delta} \left[1 + \frac{1}{(1-y)^2} \right] \sin \frac{m_{\gamma\gamma}^2 t}{4q_{(\gamma)}^0}, \\
I_2(P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) &= -\delta_{\alpha\beta} \delta_{\alpha\gamma} \sin \frac{m_{\gamma\delta} m_{\delta\gamma} t}{4q_{(\gamma)}^0}, \\
I_3(P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) &= -\delta_{\alpha\beta} \delta_{\beta\delta} \frac{1}{(1-y)^2} \sin \frac{m_{\gamma\delta} m_{\delta\gamma} t}{4q_{(\gamma)}^0};
\end{aligned} \tag{36}$$

$$\langle I_{\{W,Z\}}(m) \rangle = \frac{2}{\pi} \int_0^\pi d\theta I_{\{W,Z\}}(m, \theta) = \sum_{i=1}^3 \langle I_i \rangle \tag{37}$$

One notices that, since $q_{(\gamma)}^0$ is the incoming neutrino energy, the intensities in (36) depend on the scattering angle θ only through y (compare with (35)). So in (37) it is shown how to get averaged, over the scattering angle, the total intensity $\langle I_{\{W,Z\}}(m) \rangle$ from the sum of individual averaged intensities. Since the dimensionless intensities serve also as unnormalized probabilities, one has to look just at zero to positive values when connecting to observations; this has to be done for the individual as well as for the total intensities, unaveraged and averaged:

$$\begin{aligned}
\text{Observ..} &: I_i \geq 0, \langle I_i \rangle \geq 0 \\
\text{Unobserv.} &: I_i \leq 0, \langle I_i \rangle \leq 0
\end{aligned} \tag{38}$$

$$\begin{aligned}
\text{Observ.} & : I_{\{W,Z\}}(m, \theta) \geq 0, I_i \geq 0; \langle I_{\{W,Z\}}(m, \theta) \rangle \geq 0, \langle I_i \rangle \geq 0 \\
\text{Unobserv.} & : I_{\{W,Z\}}(m, \theta) \leq 0, I_i \geq 0; \langle I_{\{W,Z\}}(m, \theta) \rangle \leq 0, \langle I_i \rangle \geq 0 \quad (39)
\end{aligned}$$

One notices that (36) has three independent parts as the flavor numbers γ and δ are generally independent from α and β . Hence (36) is describing, respectively, the flavor conserving $\alpha = \beta = \gamma = \delta$ transition, and the flavor violating $\alpha = \beta = \gamma \neq \delta$ and $\alpha = \beta = \delta \neq \gamma$ transitions. Already at this point, one can see that one has here the physics beyond the SM. The interesting thing is that if one forces γ and δ to be identical with α one obtains that (36), (37) ≈ 0 for any value of t or L ; this is another indication that physics went beyond the SM.

At this point, it is the easiest to continue if one has the explicit values for neutrino masses and the neutrino mixing matrix. The masses accepted here are from the analysis by Fritzsch [18] with values,

$$m_1 = 0.004eV, m_2 = 0.01eV, m_3 = 0.05eV \quad (40)$$

while the neutrino mixing matrix is due to Harrison, Perkins and Scott [19]

$$(U_{\alpha i}) = \begin{pmatrix} \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & -\sqrt{\frac{1}{2}} \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \end{pmatrix} \quad (41)$$

Now, taking that the target at rest is an electron, one has $M_1 = M_2 \approx 0.5MeV$, while one retains the incoming energy of the neutrino at $q_{(\gamma)}^0 = 10MeV$ and looks for $L_+ = t_+c$, and $L_- = t_-c$, at which the values of $+\sin$ and $-\sin$ functions in (36) become $+1$, yielding first large intensities. The result is

$$\sin \frac{m_{\gamma\gamma}^2 t_+}{4q_{(\gamma)}^0} = 1 \rightarrow L_+ = \frac{2\pi q_{(\gamma)}^0}{m_{\gamma\gamma}^2}; -\sin \frac{m_{\gamma\delta} m_{\delta\gamma} t_-}{4q_{(\gamma)}^0} = 1 \rightarrow L_- = \frac{6\pi q_{(\gamma)}^0}{m_{\gamma\delta} m_{\delta\gamma}} \quad (42)$$

Next, one realizes that, to a good approximation, these distances away from the source, where the first large intensities occur, are given as $\Delta L_{\pm} = L_{\pm} - L_0 \approx L_{\pm}$, that is, with just $L_{\pm}$. As to the specific place where one notices these maxima, one has to involve the kinematics from (31) and (35).

Relation (42) is a general relation; as already mentioned, from (36), one actually has three possibilities for transitions: one flavor conserving and two flavor violating. Using the values for the neutrino masses in (40) and the neutrino matrix in (41), one deduces the distances from (42) to be

$$\begin{aligned}
& (Flav.conserv.) I_1 (P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \delta_{\alpha\beta} \delta_{\alpha\gamma} \delta_{\gamma\delta}; \\
\alpha, \beta &= e; \gamma, \delta = e : L_+ (\nu_{(e)} \rightarrow \nu_{(e)}) \approx 281 km. \\
& (Flav.viol.) I_2 (P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \delta_{\alpha\beta} \delta_{\alpha\gamma}; \\
\alpha, \beta, \gamma &= e; \delta = \mu : L_- (\nu_{(e)} \rightarrow \nu_{(\mu)}) \approx 9279 km. \\
& (Flav.viol.) I_3 (P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \delta_{\alpha\beta} \delta_{\beta\delta}; \\
\alpha, \beta, \delta &= e, \gamma = \mu : L_- (\nu_{(\mu)} \rightarrow \nu_{(e)}) \approx 9279 km.
\end{aligned} \tag{43}$$

The distances are rather large, particularly for the flavor violating oscillations, $\nu_{(e)} \rightarrow \nu_{(\mu)}$ and $\nu_{(\mu)} \rightarrow \nu_{(e)}$, which one can attribute to the small values of neutrino masses. One notices that among three distances in (43) two are the same, $L_- (\nu_{(e)} \rightarrow \nu_{(\mu)}) = L_- (\nu_{(\mu)} \rightarrow \nu_{(e)})$.

For I_1 and I_3 intensities it helps to look at θ dependence of $(1-y)^{-2}$ in order to see at which angles they are best detected. Here are some of the most significant values.

$$\begin{aligned}
& (1-y)^{-2} \left(\theta = 0; 0.1; 0.4; 0.5; \frac{\pi}{2} \right) \\
& = 1681; 60.34; 2.26; 1.7; 1
\end{aligned} \tag{44}$$

Indeed, I_1 and I_3 have rather large maxima in forward, $\theta = 0$, direction, direction that should be favorable for detecting neutrino oscillation.

One may as well look at the average values of intensities over the scattering angle θ . For this one needs the scattering angle average quantity $\langle (1-y)^{-2} \rangle$ in (36). For the electron target, which is a very light target, one notices very large values in the forward, $\theta = 0$, direction and rather smooth thereafter up to $\theta = \pi/2$. The large value in question of 1681 is at $\theta = 0$. This comes basically from the smallness of the electron mass. Another meaning of the large value at $\theta = 0$ is that the recoil activities are strong at these angles when the target is light, and weak in the perpendicular directions. The straightforward numerical integration from more detailed evaluation of $(1-y)^{-2}$ yields for the average values

$$\begin{aligned}
\langle (1-y)^{-2} \rangle &= \frac{2}{\pi} \int_0^{\pi/2} d\theta (1-y)^{-2} \approx 59, \\
\langle I_1 (P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \rangle &= \delta_{\alpha\beta} \delta_{\alpha\gamma} \delta_{\gamma\delta} 60 \sin \frac{m_{\gamma\gamma}^2 t}{4q_{(\gamma)}^0} \\
\langle I_2 (P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \rangle &= -\delta_{\alpha\beta} \delta_{\alpha\gamma} \sin \frac{m_{\gamma\delta} m_{\delta\gamma} t}{4q_{(\gamma)}^0} \\
\langle I_3 (P_{1(\alpha)} + \nu_{(\gamma)} \rightarrow P_{2(\beta)} + \nu_{(\delta)}) \rangle &= -\delta_{\alpha\beta} \delta_{\beta\delta} 59 \sin \frac{m_{\gamma\delta} m_{\delta\gamma} t}{4q_{(\gamma)}^0}
\end{aligned} \tag{45}$$

The numbers 60 ,1 and 59 have the meanings or relative strengths with respect to each other of these dimensionless intensities. Then according to definitions in (38) and (39), for flavor conserving part, $\langle I_{\{W,Z\}}(m, \theta) \rangle \approx 100\% \langle I_1(\alpha + \alpha \rightarrow \alpha + \alpha) \rangle$, where one identified $60 = 100\%$. The flavor violating parts $\langle I_2 \rangle$ and $\langle I_3 \rangle$ one may say that they split 100% as $\langle I_{\{W,Z\}}(m, \theta) \rangle \approx 2\% \langle I_2(\alpha + \alpha \rightarrow \alpha + \delta) \rangle + 98\% \langle I_3(\alpha + \gamma \rightarrow \alpha + \alpha) \rangle$.

Now, unlike in the classical field theoretical model of Dvornikov [17], here one has quantum field theory with the particle kinematics so that scattering angle θ between the incoming neutrino and scattered electron come into play. So far one was dealing with the incoming neutrino and derived the intensities. However, at the distances of $281km$ and $9279km$ one can hope to detect the outgoing neutrino energy $q_{(\delta)}^0$ while, at the same time measure the scattering angle θ of the recoiled electron. From relations (31) and (35) one now correlates $q_{(\delta)}^0$, $q_{(\gamma)}^0$ and angle θ through the relation

$$\begin{aligned} P_{(1)} &= \left(\vec{0}, M_1 \right), \vec{q}_{(\gamma)} \cdot \vec{q}_{(\delta)} = q_{(\gamma)}^0 q_{(\delta)}^0 \cos \phi : \\ & q_{(\gamma)}^{02} \sin^2 \theta \left[q_{(\gamma)}^0 - q_{(\delta)}^0 + 2M_1 \right] \\ + q_{(\gamma)}^0 M_1 \left(M_1 - 2q_{(\delta)}^0 \right) - q_{(\delta)}^0 M_1^2 &= 0; \end{aligned} \quad (46)$$

$$q_{(\gamma)}^0 = \left| \vec{P}_{(2)} \right| \cos \theta + q_{(\delta)}^0 \cos \phi, \quad (47)$$

$$\left| \vec{P}_{(2)} \right| \sin \theta = q_{(\delta)}^0 \sin \phi \quad (48)$$

Relation (46), for example, associates the incoming neutrino energy with a measured outgoing neutrino energy. Relations (47) and (48) correlate the electron recoil with incoming and outgoing neutrino energies. Specifically, with the knowledge of θ and $q_{(\gamma)}^0$, one can calculate the value for $q_{(\delta)}^0$; for instance for $\theta \approx 0$, (46) gives $q_{(\delta)}^0 \approx 0.24MeV$. From (47), one has that $|\cos \phi| = \left| q_{(\gamma)}^0 - \left| \vec{P}_{(2)} \right| \cos \theta \right| \left(q_{(\delta)}^0 \right)^{-1} \leq 1$. From (47) and/or (48), one can solve for two quantities, assuming that others are known. The presumed oscillation of the δ -neutrino is occurring along the line with the angle ϕ with respect to $\vec{q}_{(\gamma)}$ and the corresponding intensity is determined by relations (45). Although, by and large, one expects $q_{(\delta)}^0 < \left| \vec{P}_{(2)} \right|$ which implies $\sin \theta < \sin \phi$, never the less, smaller angle θ would imply also smaller angle ϕ . Of course in addition to relations (46) and (47) one can find other relations that can serve the purpose of studying the neutrino oscillations from scattering experiments.

Discussion— One thing that one notices right a way is the fact that while the LIV is very real, because it is associated with the $O(m^2)$ terms, it is negligible at least in the scattering-like experiments. Therefore, the "mass-less" SM is basically LI because the neutrinos have masses that are $\leq 1eV$. Because the LIV terms are proportional to $O(m^2)$ they play no role in the laboratory

experiments where the SM dominates. As shown by relations (30)- (34), the extrapolation of the negligible portion of the laboratory neutrino oscillation scattering cross-sections into the practical baseline oscillation differential cross-sections should make them now observable at reasonable distances from the interaction region, particularly through the help of the neutrino cross-section intensity (40). The most interesting thing is that this extrapolated differential cross-section contains the flavor conserving and two kinds of flavor violating parts.

Acknowledgements— I wish to thank Dr. Howard E. Brandt for friendly discussions and help with the computer manipulations. Thanks are due to the referee (anonymous) for pointing out to additional and necessary clarifications which extended the manuscript beyond the laboratory scattering to the baseline oscillation region. To my wife Patricia Marie Stone Soln, I am deeply grateful for her expert librarian help in locating literature over the internet.

References—

- [1] Super-KamiokandeCollaboration, Y. Ashie et al., Phys. Rev. Lett. **93**, 101801 (2004); M. Shiozawa, Prog. Part. Nucl. Phys. **57**, 79 (2006).
- [2] SNO Collaboration, Phys. Rev. Lett. **81**, 071301 (2001); **89**, 011301 (2002); **89**, 011302 (2002); Phys. Rev. **C72**, 055502 (2005).
- [3] KAMLAND Collaboration, T. Araki et al., Phys. Rev. Lett. **94**, 081801 (2005).
- [4] Homestake Collaboration, T. Leveland et al. Astrophys. J. **496**, 505 (1998); GNO Collaboration, M. Altman et al. Phys. Lett. **B616**, 174 (2005); SAGE Collaboration, J. N. Abdurashitov et al., Nucl. Phys. Proc. Suppl. **110**, 315 (2002); Super-KamiokandeCollaboration, J. Hosaka et al., Phys. Rev. **D73**, 112001 (2006).
- [5] S. M. Bilenky, C. Giunti and W. Grimus, Progress In Particle and Nuclear Physics, **43**, 1 (1999).
- [6] C. Giunti and M. Laveder, “Neutrino Mixing”; hep-ph/0310238v2.
- [7] B. Kayser, “Neutrino Oscillation Phenomenology”; Proc. of the 61 st Scottish Universities Summer School in Physics, Eds. C. Frogatt and P. Soler (to appear); arXiv: 0804.1121v3 [hep-ph].
- [8] M. Fukugita and T. Yanagida, “Physics of Neutrinos and Applications to Astrophysics” (Springer Verlag Berlin Heidelberg 2003).
- [9] C. Giunti and C. W. Kim, “Fundamentals of Neutrino Physics and Astrophysics” (Oxford University Press, Oxford 2007).
- [10] C. Giunti, “Neutrino Flavor States and the Quantum Theory of Neutrino Oscillations” (XI Mexican Workshop on Particles and Fields, 7-12 November 2007), arXiv: 0801. 0653 v1 [hep-ph].
- [11] C. Giunti, “Fock States of Flavor Neutrinos are Unphysical”; Eur. Phys. J. **C39**, 377-382 (2005); hep- ph/0312256v2.
- [12] B. Pontecorvo, JETP **7**, 172 (1958).
- [13] R. E. Schrock, Phys. Lett. **96B**, 159 (1980).

- [14] Y. F. Li and Q. Y. Liu, JHEP 0610, 048 (2006).
- [15] M. Blasone, A. Capolupo, F. Terranova and G. Vittiello, Phys. Rev. **D73**, 013003 (2005).
- [16] S. M. Bilenky, F. von Feilitzsch and W. Potzel, J. Phys. G; Nucl. Part. Phys. **34**, 987 (2007), hep-ph/0611285v2.
- [17] M. Dvornikov, Phys. Lett. **B610**, 262 (2005), hep-ph/0411101.
- [18] H. Fritzsch, "Flavor, Mixing, Neutrino Masses and Neutrino Oscillations", arXiv: 0902.2817.
- [19] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. **B530**, 167 (2002).